

柠檬烯和柠檬烯氧化物与硫酸的非均相反应动力学

王天鹤^{1,2} 刘 泽¹ 王炜罡^{1,*} 葛茂发^{1,*}

(¹中国科学院化学研究所分子动态与稳态结构国家重点实验室·北京 100190; ²中国科学院研究生院, 北京 100049)

Heterogeneous Uptake Kinetics of Limonene and Limonene Oxide by Sulfuric Acid Solutions

WANG Tian-He^{1,2} LIU Ze¹ WANG Wei-Gang^{1,*} GE Mao-Fa^{1,*}

(¹State Key Laboratory for Structural Chemistry of Unstable and Stable Species, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100190, P. R. China; ²Graduate University of Chinese Academy of Sciences, Beijing 100049, P. R. China)

*Corresponding authors. WANG Wei-Gang, Email: wangwg@iccas.ac.cn; Tel: +86-10-62558682. GE Mao-Fa, Email: gemaofa@iccas.ac.cn; Tel: +86-10-62554518; Fax: +86-10-62559373..




Fig.S1 Mass Spectrum of α -terpinene




Fig.S2 Mass Spectrum of γ -terpinene




Fig.S3 Mass Spectrum of terpinolene




Fig.S4 Mass Spectrum of 1-terpineol




Fig.S5 Mass Spectrum of α -terpineol




Fig.S6 Mass Spectrum of γ -terpineol




Fig.S7 Mass Spectrum of Peak 7 in Fig.5. The proposed corresponding molecule is 1, 4-terpin hydrate




Fig.S8 Mass Spectrum of cis-terpin hydrate




Fig.S9 Mass Spectrum of trans-terpin hydrate